

Genetic variation analysis of cytb gene of cattle bred in Iraq

¹Jaafar M. Owaid ²Asaad Y. Ayied ³Fayza A. Ahmed⁴

^{1,2,3}Animal Production Department, College of Agriculture, ⁴ Biology Department, College of Education for Pure Science, University of Basrah, Basrah, Iraq

asaad.yheia@gmail.com

Abstract

The present study was conducted at laboratory of Genetic Engineering at the University of Basrah from 01/04/2017 to 01/08/2018. The aim of study was identify genetic variations and SNPs in mtDNAcytb; among Holstein, Local Iraqi cattle and their crosses. The primer used in this study amplified 1139-bp fragments from cytb gene. The results showed the presence of 3, 12 and 2 polymorphic sites leading to the construction of 3 different haplotypes for Holstein, local and crosses respectively. Haplotype and nucleotide diversity were (0.275; 0.00047), (0.417, 0.00068), and (0.733, 0.00686) respectively. Neighbor-joining trees were constructed using 34 samples showed that all studied cattle appeared into haplotype 1(H1), while Holstein also appeared in H5 and H6 (with different two or one bases respectively). Local breed included in H2 and H3 (with one different base), the crosses in H4(with 11 different bases). AMOVA showed that variation within breed (between individuals was higher (65.35%) than between breeds (34.65%). Neutrality test both Tajima's D and Fu's Fs revealed that Holstein recorded the highest negative values (-1.67053 and -0.24213 respectively). Whereas, the crosses cattle recorded positive values (1.31709 and 6.05664 respectively). The local breed showed positive and negative values near to zero (-0.93613 and 0.01635 respectively). It can be concluded from these results that Iraqi local breeds showed high genetic variability and totally differ from other breeds in the world.

Keywords: Cattle, cytb, MtDNA, haplotype diversity, nucleotide diversity, AMOVA.

I. Introduction

Mitochondrial DNA (mtDNA) is a powerful tool that can be used to determine evolutionary relationships, population composition and biology of many species due to its low molecular weight properties, simple structure, low recombination rate and rapid evolution rate (Curole and Kocher, 1999; Wonget *al.*, 2004; Arif and Khan, 2009; Patwardhan et al., 2014; Hussain *et al.*, 2015).

The cytochrome b (Cytb) gene is 1139 base pairs and is located between nucleotide pairs 14156 and 15295(NCBI, 2007). An encoded mitochondrial gene (Cardoso et al., 2008). This gene is involved in the transmission chain of the electron in the respiratory tract of the mitochondria (Hsieh et al., 2001; Linacre and Tobe, 2011).It is used to translate protein across membranes to synthesize ATP and various cellular processes (Chen et al., 2009).It is one of the world's standard genes for comparing evolution rates (Awise, 2004). And in the studies of evolution and development and inheritance of breeds and the classification of different species(Johns and Awise, 1998;Li et al., 2005; Zhong et al., 2014). Comparative studies based on Cytb lead to the recognition of new species and breeds, as well as help in understanding evolutionary relationships due to the change in Cytbsequences(Castresana, 2001). Mitochondrial genes have been used to study genetic variation in many agricultural animals such as sheep (Othman et al., 2018; Ayied and Zqeer, 2018), goats (Amer, 2014),cattle (Tety et al., 2015) and camel (Ayied et al, 2018). The objectives of the present study were to estimate genetic variation caused by Cytb among different breeds of cattle in Iraq.

II. Materials and Methods

The study was conducted for the period from 01/04/2017 to 01/08/2018, at the laboratory of Genetic Engineering at the University of Basrah, following with the collection of data from the field up to 15/03/2018. The study included the use of 40 cows (20 Holstein, 10 local and 10 crosses). The blood samples (5ml/cow) from the jugular vein were collected. Milk chemical contents included fat%, protein%, lactose% and solid not fat (SNF%) were estimated by Lactoflash produced by Funke Gerber, Germany. Samples of milk (50 ml) were collected every 15 days throughout the experiment period.

The analyses were carried out on 40cows. Blood from each cow was sampled intravitally into sterile vacuum tubes containing K₂EDTA (dipotassium ethylene diamine tetra acetic acid) anticoagulant. A fragment (1250bp) of the mtDNACytb in the reference cattle mitochondrial genome by using the primer F-CTCCATCAACAAGCCAGTA 3'-R-TGTGTAGTAGGGGATTAGAGCA(Vakalounakis and Fragkiadakis, 1999).

The PCR amplifications were conducted in a 50 µl volume containing 20 ng genomic DNA, 25 µl of Master Mix, 2 µl each primer, 15 µl free water. The amplification conditions were as follows: initial denaturation at 94 C for 2 min followed by 35 cycles of denaturation at 94 C for 0.5 min, annealing at 58 C for 0.5min, and extension at 72 C for 0.5 min, and then the final extension at 72 C for 10 min. The PCR products were electrophoresed on 2% agarose gel stained with ethidium bromide to test the amplification success. The amplified products were purified with a DNA purification kit (SSufine) according to the manufacturer's instructions to remove residual primers and dNTPs. Sequencing was performed in sync™ DNA Extraction Kit was used for DNA extraction and manufactured by the Taiwanese Geneaid company.

Data Analysis

ATPase8/6 sequences were aligned using the BioEdit software (Hall, 1999). Haplotype diversity (HD) and nucleotide diversity (π) were analyzed using DnaSP v5. 10 software (Librado and Rozas, 2009). Genetic distance, molecular variation (AMOVA) and neutrality test were analyzed using Arlequin 3.5.1.2 software (Excoffier and Lischer, 2010). The haplotypes network was drawn using Network 5.0.0.0 software (Bandelt et al., 1999). Neighbor-joining (NJ) tree for testing cattle breed sequences and the phylogenetic tree the three genotypes (Holstein, Local and crosses) were constructed using Megaversion 7.0 software (Kumar *et al*, 2016).

Statistical analysis

A complete randomized design (CRD) was used to analyze the production data using the SPSS (2016) statistical program Version 24 and compared the means using the General Linear Model within the program. The model included two factors, the first was the breed (Holstein, local and their crosses) and the second was the haplotypes within breed.

Results and Discussion

Genetic diversity

Number of Ctyb sequences were 32 (table, 1). Ten, sixteen and six of sequences belonged to local cattle, cross and Holstein respectively. Haplotypes number (H) were nine distributed equally (3 each) on local, crosses and Holstein. Local, crosses and Holstein cattle showed a polymorphisms (NH) of 2, 13 and 3 respectively. Crosses revealed the highest value of haplotype diversity (HD) (0.733), followed by the local breed (0.417) and the Holstein cattle (0.275). Cross cattle also recorded highest nucleotide diversity (π) followed by local and Holstein cattle (0.00686, 0.00068 and 0.00047 respectively).

Table (1) Genetic diversity of Ctyb gene among different cattle breeds

Breeds	Number of Sequences (N)	Haplotype number (H)	Number of Polymorphisms (NH)	Haplotype diversity (HD)	Nucleotide Diversity (π)
Local	10	3	2	0.417	0.00068
Crosses	16	3	13	0.733	0.00686
Holstein	6	3	3	0.275	0.00047

Haplotype network

A total number of haplotypes of Cytb gene showed by different breeds were six (fig. 1). The central circle represents Haplotype 1 (H₁). Five branches appeared from H₁, the first branch was H₂ which differed from H₁ by one base (111) and represent the local cattle only. The other branch represented H₃ shown by local breed and differed from H₁ by one base (663) also. Whereas, the haplotype H₄ showed by cross cattle which varied by 11 bases from H₁. H₅ and H₆ represented the Holstein cattle and differed from H₁ by 2 and 1 bases respectively.




Fig. (1) Haplotype network of Cytb gene among studied cattle

Genetic Distance

The genetic distance of the Cytb gene (Table 2) showed that the highest distance was between the local breed and the Holstein breed (0.5287) and the lowest genetic distance between the Holstein and the cross cows (0.1788). The genetic distance between the local cattle breed and the cross was 0.2890.

Table (2) The genetic distances among studied cattle breeds in Iraq

Breeds	Local	Cross
Local	0.000	0.2890
Holstein	0.5287	0.1788

Evolutionary tree of Cytb gene between Iraqi cows and cows of the world

The results of the evolutionary tree compared to some of the countries of Cytb and the cows bred in Iraq (Fig. 2) showed that there were two main branches. The first branch included all cattle breeds of the comparative countries, with the Iraqi bred in Iraq. Iraqi origin is different from the rest of the cows in the countries in comparison to the tree of evolution.




Fig. 2 Neighboring tree of Cytb between cows in Iraq and some countries

Analysis of Molecular Variation (AMOVA)

AMOVA of Cytb gene among studied breeds of cattle resulted in between breed variation of 34.65% and within breed variation of 65.35% (table, 3).

Table (3) Analysis of Molecular Variance (AMOVA) of Cytb gene for studied breeds

Source of Variation	Degree of Freedom	Sum Squares	Variance components	Variation %
Between Breeds	2	9.734	0.44289	34.65
Within Breeds	31	21.714	0.83516	65.35
Total	33	31.448	1.27806	

Neutrality Test

The results of neutrality test (Tajima's D) of the Ctyb gene (table, 3) showed negative values for local breeds (-0.92613) and Holstein (-1.67053), but it was positive (1.21709) for cross cows. Those of Fu's Fs were 0.01635, 6.05664 and -0.24213 for local, cross and Holstein cattle respectively.

Table (3) Neutrality test of Ctyb gene for studied cattle breeds

Breeds	Tajima's test (D)	Fu's Fs test
Local	-0.93613	0.01635
Cross	1.31709	6.05664
Holstein	-1.67053	-0.24213

III. Discussion

The results of the present study showed that polymorphism of cross breed reflecting that the source of genetic variation is the crossing processes or migration. These results are in agreement with those of Dadi et al (2009) in Ethiopian cross cattle. However, local breed exhibited low polymorphism which indicated that genetic content of this breed has not changed and the main factor changing the genetic variation is the genetic drift as a result of raising this breed in very small herds (not more than 3 cows/breeder). As well as this breed distributed in remote areas and kept by breeder using very old not scientific method of breeding inherited from their ancestors. These practices elevated inbreeding which increases homozygosity and genotypic fixation. Especially it showed clear

reduction in nucleotide diversity of the studied gene and revealed clear genetic differences from both the Holstein and their crosses. Besides crossing local breed with Holstein is not under systematic crossing program with no clear target.

The results of the present study showed that the cross cows had the highest genetic makeup, the haplotypes diversity and nucleotides diversity of the Cytb gene. This indicates that the source of genetic variance in this group of animals is the process of crossing and migration. These results were in agreement with those of Dadi et al. (2009) in Ethiopian cross cattle. The Iraqi local cattle breed were characterized by the lowest genetic formation of Cytb gene, which shows that the genetic content of this breed has not changed and the main factor in changing genetic variance is genetic drift. This is the result of breeding in very small herds not exceeding three cows per breeders as well as their presence in remote areas and the holder do not have the desire to conduct breeding programs and genetic improvement or modern techniques such as artificial insemination technique, which led to an increase in the inbreeding and this in turn increased the structures and showed a clear reduction in the nucleotide diversity of Cytb gene. Local breed genetic makeup has moved away from the Holstein breed and the cross cows, especially since the cross cows in the current study may be constructed from an unprogrammed crossing local and the Holstein breeds, especially as some of them contained a high percentage of the genotypes of the Holstein blood. Tarekegn et al. (2018) found different genetic parameter of Cytb diversity, that the average haplotype diversity of Abigar and Sheko breeds were 0.8333 and 0.6282 respectively and nucleotide diversity was 0.7540 and 0.0010 respectively. The local breed was characterized by the nucleotide diversity, which was comparable to the Spanish breed Lidia cattle (Cortés et al., 2008) and higher than other European cattle whose values ranged from 0.0011-0.0057 (Loftus et al., 1994).

Cytb gene is usually used to distinguish genetic variation among animals, in a study using two local Indonesian breeds, Kebumen and Brahma, which obtained 8 and 6 haplotypes respectively and polymorphism of 36 and 7 respectively (Tarekegn et al., 2018). Cytb is a gene derived from maternal inheritance that is used to detect the origin of cattle. Romaino et al. (2014) found the possibility of distinguishing between local Malaysian cattle and wild animals using this gene as well as other studies in the world such as the Indonesian study (Sutarno and Setyawan 2015), who compared the breed of Bally and Zebo cattle.

The present study showed that there were three haplotypes of the studied breeds, which was significantly lower than Chung (2013) in the Korean and Japanese black cattle (14-10 haplotypes respectively). The number of haplotypes may vary due to sample size (Tarekegn et al., 2018). The polymorphisms of the two breeds was 29. Cytb gene polymorphism can be used to predict genetic diversity (Cai et al., 2007).

In conclusion the haplotypenetwork showed that local cattle were differentiated from the other studied breeds and recorded in the NCBI site, the local is of different origin from the rest of the international breeds, especially as Iraq is one of the first centers of domestication.

IV. References

1. **Amer, S.** (2014). Mitochondrial DNA variability among some Saudi Arabian goat breeds. *British Biotechnology Journal* 4(8): 877.
2. Arif, I. A. and Khan, H. A. (2009). Molecular markers for biodiversity analysis of wildlife animals: a brief review. *Anim. Biodivers. Conserv.* 32: (1):9-17.
3. **Avise, J. C. (2004).** Molecular markers, natural history, and evolution, 2nd edition. Sinauer Associates, Sunderland, Massachusetts
4. **Ayied; A, Y, Al-Badran; A, I, Al-Zaalan; A, R.(2018).** Assessment of Genetic Diversity in Iraqi Camel Breeds using Cytochrome b Advances in Animal and Veterinary Sciences Adv. Anim. Vet.Sci6(7):
5. **Ayied; A, Y and Zaqeer; B, F.(2018).** Polymorphism Of Coi Gene And Its Association With Milk Production And Lambs Growth Before Weaning Of Iraqi Awassi Sheep. *Int. J. Adv. Res.* 6(12),1317- 1323.
6. **Bandelt H. J., Forster, P. and Rohl, A. (1999).** Median-Joining Networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.*, 16(1): 37-48.
7. **Cai, X., Chen, H., Lei, C., Wang, S., Xue, K. & Zhang, B. (2007).** mtDNA diversity and genetic lineages of eighteen cattle breeds from Bostaurus and Bosindicus in China. *Genetica* 131(2): 175-183.
8. **Cardoso, S., González-Fernández, M., Odriozola, A., Valverde, L. & de Pancorbo, M. (2008).** Cytochrome b for species identification of biological traces found in food: A case report. *Forensic Science International: Genetics Supplement Series* 1(1): 589-590.
9. **Castresana, J. (2001).** Cytochrome b phylogeny and the taxonomy of great apes and mammals. *Molecular Biology and Evolution* 18(4): 465-471.
10. **Chen, J., Bai, Z., Gao, C. & Wang, J. (2009)a.** Morphology of Rhinencephalon and Hippocampal formation of the Bactrian Camel (*Camelus bactrianus*) with their adaptive features. *Veterinary research communications* 33(1): 25-32
11. **Chung, H. (2013).** Phylogenetic analysis and characterization of mitochondrial DNA for Korean native cattle. *Open Journal of Genetics* 3(01): 12.
12. **Cortés, O., Tupac-Yupanqui, I., Dunner, S., García-Atance, M., García, D., Fernández, J. & Cañón, J. (2008).** Ancestral matrilineages and mitochondrial DNA diversity of the Lidia cattle breed. *Animal Genetics* 39(6): 649-654.
13. **Curole J. P. and Kocher T. D. (1999).** Mitogenomics: digging deeper with complete mitochondrial genomes. *Trends in Ecology & Evolution* 14(10):394-398. DOI: 10.1016/S0169-5347(99)01660-2

14. **Dadi, H., Tibbo, M., Takahashi, Y., Nomura, K., Hanada, H. and Amano, T. (2009).** Variation in mitochondrial DNA and maternal genetic ancestry of Ethiopian cattle populations. *Animal genetics* 40(4): 556-559.
15. **Excoffier L. and Lischer H. E. (2010).** Arlequin suite ver. 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10 (3): 564-567.
16. **Hall T. A. (1999).** BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium*, 41: 95–98.
17. **Hsieh, H.-M., Chiang, H.-L., Tsai, L.-C., Lai, S.-Y., Huang, N.-E., Linacre, A. & Lee, J. C.-I. (2001).** Cytochrome b gene for species identification of the conservation animals. *Forensic Science International* 122(1): 7-18.
18. **Hussain, T., Babar, M., Musthafa, M., Saif, R., Hussain, F., Aqeel, M., Naveed, N., Pervez, M., Khan, W. and Ziaullah, S. (2015).** Mitochondrial ATP6 and ATP8 genes based molecular diversity and phylogenetic analysis in Punjab urial (*Ovis montanus*). *J. Anim. Plant Sci.* 25(3): 311-318.
19. **Johns, G. C. & Avise, J. C. (1998).** A comparative summary of genetic distances in the vertebrates from the mitochondrial cytochrome b gene. *Molecular Biology and Evolution* 15(11): 1481-1490.
20. **Kumar S, Stecher, G and Tamura K, (2016).** MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33 (7): 1870-1874.
21. **Li, A., Zhao, Q., Tang, S., Zhang, Z., Pan, S. & Shen, G. (2005).** Molecular phylogeny of the domesticated silkworm, *Bombyx mori*, based on the sequences of mitochondrial cytochrome b genes. *Journal of Genetics* 84(2): 137-142.
22. **Librado P and Rozas J, (2009).** DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25 (11): 1451–1452.
23. **Linacre, A. & Tobe, S. S. (2011).** An overview to the investigative approach to species testing in wildlife forensic science. *Investigative genetics* 2(1): 2.
24. **Loftus, R. T., MacHugh, D. E., Bradley, D. G., Sharp, P. M. & Cunningham, P. (1994).** Evidence for two independent domestications of cattle. *Proceedings of the National Academy of Sciences* 91(7): 2757-2761.
25. **NCBI, (2007).** https://www.ncbi.nlm.nih.gov/nuccore/NC_009849.1
26. **Othman, O. E., Germot, A., Khodary, M. G., Petit, D. & Maftah, A. (2018).** Cytochrome b Diversity and Phylogeny of Six Egyptian Sheep Breeds. *Annual Research & Review in Biology* 22(4): 1-11.
27. **Patwardhan, A., Ray, S. and Roy, A. (2014).** Molecular markers in phylogenetic studies a review. *J. Phylogenet. Evol. Biol.* 2:131-140.
28. **Romaino, S., Fazly-Ann, Z., Loo, S., Hafiz, M., Hafiz, M., Iswadi, M., Kashiani, P., Rosli, M., Syed-Shabthar, S. & Md-Zain, B. (2014).** Species identification of Malayan Gaur, Kedah-Kelantan and Bali cattle

- using polymerase chain reaction-restricted fragment length polymorphism. *Genet. Mol. Res* 13(1): 406-414.
29. **SPSS(2016)**. Statistical Package for Social Science, User's Guide for statistics Version 24, Copyright IBM, SPSS Inc., USA.
30. **Sutarno& A. D. Setyawan.** (2015). Review: Genetic diversity of local and exotic cattle and their crossbreeding impact on the quality of Indonesian cattle. *Biodiversitas*:16:327-354.
31. **Tarekegn, G. M., Ji, X.-y., Bai, X., Liu, B., Zhang, W., Birungi, J., Djikeng, A. & Tesfaye, K. (2018)**. Variations in mitochondrial cytochrome b region among Ethiopian indigenous cattle populations assert *Bostaurus* maternal origin and historical dynamics. *Asian-Australasian journal of animal sciences*.
32. **Tety, H., Putra, W., Putra, W., Volkandari, S., Volkandari, S. & Sumadi, S. (2015)**. Polymorphism of mtDNA Cytochrome b Gene of Local Cattle in Indonesia. *International Journal Sustainable Future for Human Security J-Sustain* 3(1): 21-24.
33. **Vakalounakis, D. J. and Fragkiadakis, G. A. (1999)**. Genetic diversity of *Fusariumoxysporum* isolates from cucumber: Differentiation by pathogenicity, vegetative compatibility, and RAPD fingerprinting. *Phytopathology*. 89: 161-168.
34. **Wong, B., Keogh, J. S. and Jennions, M. (2004)**. Mate recognition in a freshwater fish: geographical distance, genetic differentiation, and variation in female preference for local over foreign males. *Journal of Evolutionary Biology* 17(3): 701-708.
35. **Zhong, X., Wang, N., Hu, D., Wang, J., Liu, T., Gu, X., Wang, S., Peng, X. & Yang, G. (2014)**. Sequence analysis of cytb gene in *Echinococcusgranulosus* from western China. *The Korean journal of parasitology* 52(2): 205.